

# PRUNING CRAPEMYRTLES

We often use Crapemyrtles in the landscape because they bloom all summer long. We also value them for their peeling bark, fall color and the grace of their natural form. It is as tough as it is beautiful.

The practice of chopping off the tops of Crapemyrtle has become very commonplace. Many people believe that it is required to promote flowering; some prune because the plant is too large for the space provided; others see their neighbors doing it and feel the need to follow suit. There are



Butchering Crapemyrtles is an unnecessary and often harmful practice

some instances in which heavy pruning is necessary, but light pruning is usually all that is needed. The type and amount of pruning depends on the desired shape and size of the plant.



Prune off seedhead above a lateral bud

Crapemyrtle can be a low-maintenance plant, and the best way to ensure this is to choose the cultivar that best suits your landscape needs

before planting. There are many new cultivars in different sizes and colors. The dwarf (3 to 6 feet) and semi-dwarf (7 to 15 feet) selections now available make it easy to choose the right size plant for a certain space.

If careful consideration is given to the projected size of the mature plant, a selection can be found that will not outgrow its boundaries and can be allowed to display its graceful beauty with minimal pruning.

Crapemyrtles do not require heavy pruning to promote bloom. Flowers are produced on new growth. It will produce flowers without any pruning, although it will produce larger flowers and bloom more profusely if at least lightly pruned. Pruning in late winter or early spring will stimulate vigorous new growth in the spring. Encourage a second bloom in summer by pruning flowers immediately after they fade.

This plant prefers hot, sunny climates and in South Carolina many varieties

will grow to tree-size proportions. It is important that tree types are sited where they have a large area to spread. When given an ideal location, these tree types should be allowed to develop their natural style without whacking off their tops.

To develop a tree shape, remove all limbs growing from ground level except the three to five strongest limbs. As the tree matures, remove lower, lateral branches ("limbing up") one-third to halfway up the height of the plant. Remove branches that are crossing

or rubbing against each other and shoots growing into the center of the canopy. Make your cuts to a side branch or close to the trunk. Head back wayward and unbranched limbs.

As the tree grows taller, remove lower branches as needed. Remove any future growth from the ground to retain the desirable trunk structure. This basal sprouting may occur whether the tree is being pruned or not. Pull these out when succulent instead of pruning them.

You may feel the need to improve the appearance by removing the seed heads in late winter or early spring before growth begins. This is recommended only if they are within reach. Once this becomes a tall, mature plant, allow nature to take its course - the seedheads will drop, the plant will bloom, and the natural grace of the plant has been retained.



An unpruned tree- notice the multitude of basal shoots and the crowded canopy



The same Crapemyrtle after pruning- notice how the trunk character has been exposed and the canopy has been thinned

# CORNELIUS NURSERY

2233 South Voss Road, 713-292-0898  
1200 North Dairy Ashford, 281-493-0550

Web: [www.corneliusnurseries.com](http://www.corneliusnurseries.com)

# 523

2003